

Asian Journal of Soil Science and Plant Nutrition

Volume 10, Issue 4, Page 604-614, 2024; Article no.AJSSPN.127728 ISSN: 2456-9682

Effect of Different Temperatures on Reactions of Potassium Adsorption and Availability in Saline Soils within Nineveh Governorate, Iraq

Haneen H. A. Al-Jubory ^a and Qahtan D. Essa Al-Khafagi ^{b*}

^a Center for Arid Farming and Conservation Agriculture Research, Mosul University, Iraq. ^b Soil Science & Water Resources Department, College of Agriculture and Forestry, Mosul University, Iraq.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: https://doi.org/10.9734/ajsspn/2024/v10i4432

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: https://www.sdiarticle5.com/review-history/127728

Original Research Article

Received: 28/09/2024 Accepted: 30/11/2024 Published: 09/12/2024

ABSTRACT

Thermodynamic criteria were used to evaluate the adsorption and release of potassium in saline soils. The study was conducted on three sites that differ in salt content in Nineveh Governorate - Iraq, and each site is divided into two depths (surface and subsurface). The study was conducted at three temperatures (278, 298, 318 Kelvin). The results of the study showed that there was an increase in the amount of potassium adsorbed and the percentage of adsorption with an increase in the amount of potassium added in all the studied soil samples. For the ability of the soil to the adsorption, the results showed a sudden increase in the adsorption of potassium at the application of the high levels (1 and 2 mmolL⁻¹) of potassium in all the studied soils. Also, the highest value of

*Corresponding author: Email: qahtan_darwish@uomosul.edu.iq;

Cite as: Al-Jubory, Haneen H. A., and Qahtan D. Essa Al-Khafagi. 2024. "Effect of Different Temperatures on Reactions of Potassium Adsorption and Availability in Saline Soils Within Nineveh Governorate, Iraq". Asian Journal of Soil Science and Plant Nutrition 10 (4):604-14. https://doi.org/10.9734/ajsspn/2024/v10i4432.

the ionic activity of potassium (AR_k) was (14.05 × 10⁻³ molL^{-1/2}), at a temperature of 298 K in the subsurface soil layer of the first site, while the lowest one was (1.61 × 10⁻³ molL^{-1/2}), in the same layer of the first site at a temperature of 278 K. The values of free energy of potassium, ΔG_{K} , in the studied soils ranged from the highest value reached (-14.85×10⁻³ kJ) at a temperature of 278 K, to the lowest value, reached (-10.56×10⁻³ KJ) at a temperature of 298 K, in the subsurface soil layer of the first site. Regarding the values of entropy reaction, they ranged from the highest value of -53.43 Jmol⁻¹ in the subsurface depth of the first site at a temperature of 278 K. On the lowest value of -35.46 in the same layer of the first site at a temperature of 298 K. On the other hand, the enthalpy values of the studied soil samples were negative, which indicates that the reaction is of the exothermic type, where the highest negative value for enthalpy was -29.70 kJmol⁻¹ at a temperature of 278 K in the subsurface soil layer of the first site at a temperature of 298 K.

Keywords: lonic activity of potassium; free energy of potassium; entropy of potassium reaction; enthalpy of potassium.

1. INTRODUCTION

The process of potassium adsorption into the soil system is a chemical process that affects the available content of potassium. This process depends on the state of balance between the potassium present in the soil solution and the potassium that is located between the inner layers of minerals. The stabilized potassium constitutes 5-10% of the total potassium content in Soil [1]. Mouhamed et al. [2] indicated that the potassium adsorption process is affected by several factors, including soil aeration, soil moisture content, soil temperature, and quality of the prevailing clay minerals, and that the amount of available potassium constitutes approximately 1-2% of the total amount of potassium in the soil. Soil temperature has a role in the available and movement of potassium in the soil system, as well as its role in the process of its absorption by the plant. In general, the speed of any chemical reaction increases with increasing temperatures whenever the temperature increases by 10 degrees Celsius. Al-Obaidi and Al-Zubaidi [3] see the necessity of studying potassium reactions under the influence of different temperatures, especially in Iraqi environmental conditions, due to the large temperature fluctuations during the seasons of the year and during the day and night in a single day, in their study on the adsorption of potassium in calcareous soil from northern Iraq at three absolute temperatures (278, 298, 318° kelvin) using mixed displacement techniques, they found that there is a highly significant effect of temperature on the potassium adsorption and release reactions and the state of equilibrium between them. Al-Obaidi et al., [4] explained that the release of potassium in the soil results from the decomposition of potassium-bearing minerals and an exchange reaction, which is very slow,

potassium in the soil solution is constantly renewed by being released from the exchanged form and prepared from the non-exchanged form. As for Iraqi soils, as is the case in most soils in arid and semi-arid regions, they have a high reserve of potassium, but the speed of its release is slow and is not sufficient to meet the need of agricultural crops for this element [5]. The process of releasing potassium is an endothermic reaction, while the process of adsorption of potassium is an exothermic reaction. High soil temperature has a significant impact on the ability of minerals to fix potassium, which affects its available for plants [6], and Najafi-Ghiri et al., [7] indicated that high temperature in calcareous soil leads to an increase in the amount of soluble potassium and exchangeable potassium, thus increasing the rate of potassium release. The ability to release potassium from the soil may be limited due to the change in the structure of potassium-bearing soil minerals and the decrease in the diffusion of potassium ions in the interlayer spaces of the minerals. The use of thermodynamic criteria in evaluating soil fertility to express potassium status and behavior in the soil helps in revealing an appropriate management method for each soil [8,9].

2. MATERIALS AND METHODS

2.1 Choosing Study Sites

The study included three sites in the Rabia area within Nineveh Governorate - Iraq, for saline soils with different salinity levels. Soil samples were collected from the surface depth (0 - 25 cm) and the subsurface depth (25 - 50 cm) to conduct routine analyses. The soil samples were air-dried and ground using a wooden hammer,

Depth	рΗ	EC	CaCO₃	O.M	CEC	Soil	separate	es (%)	Texture
(cm)		dS.m ⁻¹	gm.kg ⁻¹		Cmol.kg ⁻¹	Sand	Silt	Clay	-
First Site									
0 - 25	7.3	2.70	225	15.98	29.56	15.05	54.25	30.70	SCL
25 - 50	7.4	5.10	220	18.36	32.17	9.70	44.50	45.80	SC
Second S	Site								
0 - 25	7.2	3.00	220	11.90	29.56	34.70	34.50	30.80	CL
25 - 50	7.3	6.00	190	12.24	29.56	31.40	38.50	30.10	CL
Third Site	;								
0 - 25	7.4	7.80	195	15.98	26.95	16.50	31.00	52.50	С
25 - 50	7.6	7.80	225	13.94	26.08	11.50	33.50	55.00	С
		SCL:	Silt Clay Lo	am, SC: Si	lt Clay, CL: Cla	y Loam, C	C: Clay		

Table 1. Some physiochemical properties of the studied soils

2 mm sigure and the amount approaced by (1)

then sieved with a 2 mm sieve, and the amount of the three separations - clay, silt, sand, total calcium carbonate, organic matter, exchange capacity of positive ions, pH, and electrical conductivity (EC) - were estimated using the methods described [10]. As shown in Table 1.

2.2 Potassium Adsorption According to the Concept of Thermodynamics

Transparent plastic cans of 100 ml with a tight seal were used to study this adsorption. The cans were prepared with the number of concentrations of potassium prepared from potassium chloride (KCI) added as follows (0, 0.1, 0.2, 0.4, 0.6, 0.8, 1, 2) mmol.I⁻¹, 2.5 grams of study soil was placed in each box, and it was added these concentrations are in the form of volume and the volume is supplemented to 50 ml with 0.01 M CaCl₂ solution, after that, the plastic cans were shaken with a shaker for two hours and left to equilibrate for 48 hours at three temperatures (278, 298, 318 Kelvin), then the suspension in the cans was filtered to obtain the equilibrium solution in which potassium, calcium, magnesium, pH, and EC were determined. The amount of potassium adsorbed on the surface of the solid phase, K - quantity, was calculated, as well as the intensity of potassium in the equilibrium solution, K - intensity, which is expressed as the relative effectiveness of potassium in the equilibrium solution according to the ratio law, as mentioned by [11], and it was calculated from the following equation:

$$\mathsf{AR}_\mathsf{K} = \frac{a\,\mathsf{K}}{a\,\mathsf{Ca} + a\,\mathsf{Mg}}$$

ARK: relative effectiveness of potassium

The mathematical relationship was drawn between the amount of potassium adsorbed,

expressed by $(\pm \Delta K)$, which represents potassium in the solid soil phase (Q) quantity on the y-axis, and the intensity of potassium in the liquid soil phase (I) intensity on the x-axis, which is (AR_K), and through this Relationship, the following constants can be found for potassium:

- 1. Potassium effectiveness ratios at equilibrium (AR_κ): it represents the intersection of Q/I with the x-axis
- Free energy ∆G: it can be calculated from the equation proposed by (Woodruff, 1955):

 $-\Delta G = 2.303 \text{ RT} \log AR_{\kappa}$

- 3. Amount of potassium adsorbed on the exchange surface was calculated from the following relationship: $K.ad = (K_{in} K_{fin}) \times V/S$ K.ad: adsorbed amount,
 - Kin: added-K.

K_{fin}: balanced-K.

V: volume of solution,

S: soil mass.

- Adsorbed potassium was calculated from the following relationship: adsorbed-K = (K_{in} – K_{fin})
- Percentage of adsorption or release was calculated from the following relationship: % adsorbed or release = (adsorbed-K / Added-K) * 100
- 6. Entropy was calculated by dividing free energy by temperature

3. RESULTS AND DISCUSSION

3.1 Effect of the Added Potassium Level on Potassium Adsorption

In order to know the behavior of ion exchange of potassium in the soil of the study, the adsorption of potassium was studied, which is one of the most important chemical reactions that occur between the liquid phase and the solid phase of the soil. Through it, potassium is transferred mutually with other ions between the two phases and ends after a certain period of time when a dynamic equilibrium of the ions exchanged between the two phases occurs. Adsorption is also affected by the surface area of the mineral colloids. dissolution, temperature, reaction degree pH, and salinity [12], as Ogwada and Sparks [13] see, potassium added to the soil binds to the surface of the solid phase of the colloidal fraction in a stabilizing manner that depends on two main mechanisms:

first mechanism: involves stabilizing potassium between the layers of crystals in three-layer minerals (1:2). This type of stabilization is accompanied by the transformation of vermiculite into a mica mineral. formation of these minerals can be confirmed through the use of X-ray diffraction (XRD) and the plane (001), where the crystal dimension shrinks from (15 Å to 12.5 Å and to 10 Å). Such changes may cause a decrease in the exchange capacity of positive ions.

second mechanism: In which potassium is stabilized through the formation of insoluble potassium compounds such as aluminosilicates, and high pH values accelerate the formation of these components.

Tables (2, 3, and 4) show the amounts of potassium adsorbed, which varied according to the level of added potassium and according to the soil properties that have a direct relationship with adsorption. In general, there was an increase in the amount of potassium adsorbed and the percentage of adsorption with an increase in the amount of potassium added to the soil and for all soils of the study, the reason for this is that the specialized adsorption sites were occupied by the added potassium, which could occur at the beginning of the reaction,

 Table 2. Quantities of added and adsorbed potassium and the rate of adsorption or release to the first site at different temperatures

K	Dep	K	K	Adsorb/	Dep	K	K	Adsorb/
mmol I ⁻¹	-Ħ	mmol I ⁻¹	mmol I ⁻¹	Kelease	-Ť	mmol I ⁻¹	mmol I ⁻¹	Kelease
at 278 Kg	lvin		minol.i	/0		mmon.	1111101.1	/0
0	Sivin O	0.12	-0.12		S	0.10	-0.1	
01	urf	0.12	-0.05	-500.0	du	0.10	-0.02	-20.0
0.1	ace	0.13	0.03	15.0	ဂ	0.12	0.02	15.0
0.4) e	0.23	0.00	42.5	. .	0.23	0.00	42.5
0.6	2 N	0.30	0.3	50.0	(25	0.28	0.32	53.3
0.8	5	0.35	0.45	56.2	сл U	0.35	0.45	56.2
1	crr	0.43	0.57	57.0	9	0.43	0.57	57.0
2	2	0.76	1.24	62.0	m	0.76	1.24	62.0
at 298 Ke	elvin							
0	ဂ	0.30	-0.3		ဂ	0.30	-0.3	
0.1	Infe	0.35	-0.25	-250.0	<u>p</u>	0.35	-0.25	-250.0
0.2	lCe	0.40	-0.2	-100.0	Su	0.43	-0.23	-115.0
0.4	ô	0.53	-0.13	-32.5	г. ()	0.56	-0.16	-40
0.6	-25	0.66	-0.06	-10	25	0.71	-0.11	-18.33
0.8	<u>0</u>	0.81	-0.01	-1.25	-50	0.84	-0.04	-5.0
1	ă	0.97	0.03	3.0	0	0.97	0.03	3.0
2		1.79	0.21	10.5	ä	1.84	0.16	8.0
at 318 Ke	elvin							
0	Su	0.30	-0.3		Sul	0.28	-0.28	
0.1	rfa	0.40	-0.3	-300.0	6	0.40	-0.3	-300.0
0.2	ce	0.43	-0.23	-115.0	ùr	0.43	-0.23	-115.0
0.4	(ò	053	-0.13	-32.5	โ	0.56	-0.16	-40.0
0.6	.25	0.66	-0.06	-10.0	ÿ	0.69	-0.09	-15.0
0.8	<u> </u>	0.79	0.01	1.25	50	0.79	0.01	1.25
1	ä	0.89	0.11	11.0) 0	0.86	0.14	14.0
2		1.57	0.43	21.5	т	1.66	0.34	17.0

(+): adsorption (-): release

and in the subsequent stages. Saturation of nonspecialized sites, so any amount of potassium added to the soil is offset by an increase in the amount of potassium in the equilibrium solution [14].

We also note from the same tables that the amount of potassium adsorbed and the percentage of adsorption differed from one soil to another for the soils of the study, and the reason for this variation is that potassium is affected by the processes of adsorption and sedimentation, as well as the exchange capacity of positive ions, the degree of soil reaction, the amount of calcium carbonate and organic matter, and the type and content of clay and clay minerals, potassium content and the chemical composition of the soil solution [15]. The results that we obtained included a sudden increase in potassium adsorption at a high addition rate for the added potassium level, especially at the level of (1 and 2) mmol.L⁻¹ for all study soils, the reason for this

is due to the possibility of the complete transformation of the smectite mineral into octahedral and the occurrence of a Rotation phenomenon of the hydroxyl group OH, which leads to a sudden increase in stabilization [16].

There are also three specialized sites on the surfaces of some minerals, including illite, in which the potassium binding strength varies: the surfaces, the edges, and between the layers, which makes the speed of potassium release and adsorption vary. The total charge density and its distribution between the octahedral and tetrahedral sheets greatly affects the presence of the potassium ion in the group minerals (2:1) [17]. In this regard, some minerals have the ability to re-fix the added potassium in nonexchangeable forms. Therefore, the process of releasing potassium from these minerals into forms that are soluble, exchangeable, and adsorption are reversible equilibrium reactions controlled by the energy of binding potassium

Table 3. Quantities of added and adsorbed potassium and the rate of adsorption or release to
the second site at different temperatures

K	De	K	K	Adsorb/	De	К	Κ	Adsorb/
Added	<u>pt</u>	equilibrium	adsorbed	Release	<u>pt</u>	equilibrium	Adsorbed	Release
mmol.l ⁻¹	۔	mmol.l ⁻¹	mmol.l ⁻¹	%	ر	mmol.l ⁻¹	mmol.l ⁻¹	%
at 278 Kel	vin							
0	Su	0.30	-0.3		Su	0.12	-0.12	
0.1	ŕа	0.38	-0.28	-280.0	ъ	0.17	-0.07	-70.0
0.2	Се	0.43	-0.23	-115.0	ů	0.23	-0.03	-150.0
0.4	Ô	0.51	-0.11	-27.5	$\widehat{\mathbf{n}}$	0.30	0.10	25.0
0.6	-25	0.63	-0.03	-5.0	5	0.43	0.17	28.3
0.8	<u>)</u>	0.69	0.11	13.7	50	0.51	0.29	36.2
1	Ä	0.81	0.19	19.0) c	0.51	0.49	49.0
2		1.40	0.60	30.0	В	0.86	1.14	57.0
at 298 Kel	vin							
0	Su	0.35	-0.35		Su	0.20	-0.2	
0.1	rfa	0.43	-0.33	-330.0	ъ	0.25	-0.15	-150.0
0.2	се	0.56	-0.36	-180.0	ûr	0.30	-0.1	-50.0
0.4	(0.63	-0.23	-57.5	$\widehat{\mathbf{N}}$	0.38	0.02	5.0
0.6	Ż	0.79	-0.19	-31.6	Ч	0.58	0.02	3.3
0.8	<u> </u>	0.92	-0.12	-15.0	50	0.66	0.14	17.5
1	Ä	1.04	-0.04	-4.0) c	0.74	0.26	26.0
2		1.79	0.21	10.5	В	1.30	0.7	35.0
at 318 Kel	vin							
0	S L	0.38	-0.38		Sul	0.20	-0.2	
0.1	rfa	0.46	-0.36	-360.0	5	0.28	-0.18	-180.0
0.2	ce	0.48	-0.28	-140.0	ŭŕ	0.33	-0.13	-65.0
0.4	(0.58	-0.18	-45.0	โ	0.40	0	0.0
0.6	No	0.71	-0.11	-18.3	5	0.51	0.09	15.0
0.8	<u> </u>	0.81	-0.01	-1.2	50	0.61	0.19	23.7
1	Ϊ	0.89	0.11	11.0) c	0.63	0.37	37.0
2		1.54	0.46	23.0	В	1.27	0.73	36.5

(+): adsorption (-): release

to the exchange surfaces or its presence within the interstitial spaces of the layers clay minerals 2:1 as described by [18]. Accordingly, we find that potassium that is released quickly is bound to the exchange surfaces (clay surfaces) and is controlled by the Vanderwaals bond, while potassium that is released slowly is bound to sites within the structural structure of the mineral, and its release from those sites requires more energy [19].

3.2 Ability of Soil to Absorb Potassium at Different Temperature

The results shown in Tables (2, 3, and 4) indicate that the amount of potassium fixed in the study soil samples at the different levels of added potassium was characterized by the release of amounts of potassium to varying degrees between the soils and the different temperatures. In the study soil samples, adsorption appeared at the third and fourth addition levels (0.2 and 0.4 mmol.I⁻¹) at a temperature of 278°K for both depths, except for the soil of the superficial depth of the second site, which was adsorption appeared at the sixth level of addition (0.8 mmol.I⁻¹). While at a temperature of 298°K, adsorption appeared at the seventh and eighth addition levels (1 and 2 mmol.l⁻¹) and for both depths, except for the soil at the subsurface depth of the second site and both depths of the third site, where adsorption appeared at the fourth addition level (0.4 mmol.l-1), but at the high temperature of 318°K, potassium adsorption appeared at the seventh and eighth addition levels (1 and 2 mmol.I-1) for all soils at the superficial depth. As for the subsurface depth, adsorption appeared at the fifth and sixth addition levels (0.6 and 0.8 mmol.l⁻¹). In general, we noticed a decrease in the ability of the study soil to absorb potassium at a temperature of 298°K, which represents the spring and fall temperature, and it is the best temperature at which larger amounts of potassium are released compared to the summer temperature (318°K) and the winter temperature (278°K). The reason for this may be due to rapid saturation of the specialized and non-specialized sites for potassium adsorption, which can occur in the first hours of the equilibrium period and thus occupies the space of most adsorption sites in the soil and their percentage decreases. Thus, any increase in added potassium will reduce the potassium present in the equilibrium solution and thus its spread to within the crystalline structure of clay minerals. Also, the variation in stabilization behavior with the increase of potassium added to

the soil and the difference in its ability to adsorption and release clearly reflects the role of the chemical, mineral and physical properties of these soils, and as Al-Jubouri [20] showed that there are several processes involved in the processes of release and adsorption of potassium from the soil, including those related to mineral and organic components in the soil. Since the Iragi soil is dominated by conditions of drought and salt accumulation, and is rich in minerals containing potassium, it has a decent store of this element, but with a low speed of release depending on the mineral composition of the soil, especially the quality of surface-active clay minerals that are not only important in the processes of fixation and release of potassium, but they also affect the freedom and movement of potassium in the soil through the adsorption of potassium ions on adsorption sites that differ in the extent her preference. Li et al., [21] and Das et al., [22] showed that the ability of soil to adsorb and release ions, including potassium, depends on the structural structure of the clav minerals due to the element atoms they contain within the layers of the mineral (the tetrahedral and octahedral layers). For example, it leads The presence of potassium-bearing minerals such as mica and illite significantly increases the concentration of potassium in the soil solution.

3.3 Effect of Temperature on the lonic Activity Ratios of Potassium at Equilibrium (AR⁰_κ)

lonic activity has been used in research to evaluate the available of some nutrients, including potassium, and for the purpose of identifying locations where adsorption occurs on the surfaces of the solid phase of soil, we have relied on relative effectiveness values. The values of the relative effectiveness of potassium. AR^{0} _K, represent the intensity of potassium in the solid phase of the soil when there is no loss or gain of potassium in the soil system. They are a measure of the instantaneous available content of the soil and have no relationship to the soils ability to release or prepare the ion over a long period of time [23]. The lowest value of ionic activity was recorded at the subsurface depth of the first site (1.61846×10-3) mol.I-1/2, while the highest value of ionic activity rates at the surface depth of the second site was (6.270×10⁻³) mol.l⁻ 1/2, at a temperature of 278° K and a rate of (3.20×10^{-3}) mol.l^{-1/2}, when the equilibrium temperature was increased to 298° K, it led to a high increase in the ionic activity values. We noticed that the lowest value was at the

subsurface depth of the third site (3.96×10^{-3}) mol.I^{-1/2}, and the highest value appeared at the surface depth of the site. The first is (14.0517×10^{-3}) mol.l^{-1/2}, with a rate of (8.63×10^{-3}) mol.I-1/2. As for the temperature of 318° K, we noticed that the lowest values were in the subsurface depth of the fourth site (3.6787×10⁻³) mol.l^{-1/2}. and the highest values were (10.2284×10-3) mol.I-1/2, at the surface depth of the first site, at a rate of (7.37×10-3) mol.I-1/2 (Table 5). These values indicate that adsorption occurs in locations between layers of minerals as well as at the edges, as indicated by Hamed [24] that values that are more than (1×10⁻²) mol.I^{-1/2} indicate the dominance of adsorption over (Positions). Planner), while values less than (1×10⁻²) mol.I^{-1/2} indicate the dominance of adsorption (edge site). This confirms that the process accompanied adsorption is bv stabilization processes that are slow and occur

between lavers of minerals clav is the result of rapprochement between ions and the exchange surface in soils [25]. We note that a rise in temperature above 278°K generally led to a clear increase in the relative effectiveness values, this effect of temperature was clear at the temperature of 298° K. This confirms the assumption that the adsorption process with these soils does not include superficial adsorption on the surfaces of particles only the soil, but this is accompanied by slow processes of fixing potassium between the layers of clay minerals. In this regard, Strauss and Brummer [26] indicated that the reaction of potassium with soil components includes an initial adsorption reaction, and this reaction is completed within minutes of its beginning and is then followed by a slower reaction. Increasing the temperature will increase the rate of reaction speed, which follows the surface adsorption process.

 Table 4. Quantities of added and adsorbed potassium and the rate of adsorption or release to the third site at different temperatures

K Added	Dept	K equilibrium	K adsorbed	Adsorb/ Release	Dept	K equilibrium	K adsorbed	Adsorb/ Release
mmol.I ⁻¹	-2	mmol.l ⁻¹	mmol.I ⁻¹	%	-2	mmol.l ⁻¹	mmol.l ⁻¹	%
at 278 Ke	elvin							
0	Su	0.20	-0.2		Su	0.17	-0.17	
0.1	rfa	0.23	-0.13	-130.0	ъ	0.20	-0.1	-100.0
0.2	се	0.28	-0.08	-40.0	ů	0.25	-0.05	-25.0
0.4	(o	0.35	0.05	12.5	$\widehat{\mathbb{N}}$	0.30	0.1	25.0
0.6	25	0.40	0.2	33.3	ហ៊	0.38	0.22	36.6
0.8) ପ	0.51	0.29	36.2	50)	0.73	0.07	8.7
1	З	0.56	0.44	44.0	Q	0.92	0.08	8.0
2		1.38	0.62	31.0	л	1.66	0.34	17.0
at 298 Ke	elvin							
0	Su	0.28	-0.28		Sul	0.23	-0.23	
0.1	rfa	0.33	-0.23	-230.0	0	0.28	-0.18	-180.0
0.2	ce	0.35	-0.15	75.0	ůr.	0.30	-0.1	-50.0
0.4	`	0.40	0	0.0	N N	0.35	0.05	12.5
0.6	25	0.51	0.09	15.0	ហ៊	0.46	0.14	23.3
0.8) C	0.56	0.24	30.0	00	0.56	0.24	30.0
1	Ц	0.63	0.37	37.0	5	0.63	0.37	37.0
2		1.27	0.73	36.5		1.30	0.7	35.0
at 318 Ke	elvin							
0	Su	0.33	-0.33		Sul	0.28	-0.28	
0.1	fa	0.38	-0.28	-280.0	0 0	0.30	-0.2	-200
0.2	e	0.43	-0.23	-115.0	ůr.	0.35	-0.15	-75.0
0.4	(0.48	-0.08	-20.0	Ñ	0.43	-0.03	-7.5
0.6	25	0.53	0.07	11.6	ο ά	0.51	0.09	15.0
0.8) Cl	0.63	0.17	21.2	<u>50</u>	0.58	0.22	27.5
1	н	0.71	0.29	29.0	9	0.69	0.31	31.0
2		1.22	0.78	39.0	Ц	1.15	0.85	42.5

(+): adsorption (-): release

Depth (cm)	$AR^{0}_{K} \times (10^{-3})$ mole.L ^{-1/2} at temperature				
	278° Kelvin	298° Kelvin	318° Kelvin		
First Site					
Surface (0-25)	2.02037	12.6901	10.2284		
Sub Sur. (25-50)	1.61846	14.0517	10.0280		
Second Site					
Surface (0-25)	6.2700	11.9736	8.3204		
Sub Sur. (25-50)	2.3053	4.0018	3.6787		
Third Site					
Surface (0-25)	3.0061	5.1187	8.3204		
Sub Sur. (25-50)	4.0020	3.9602	3.6787		

Table 5. Value of activity relative at equilibrium (AR^{0}_{K}) for three temperatures

3.4 Effect of Temperature on Free Energy of Potassium (ΔG)

The free energy ΔG is one of the thermodynamic functions used to describe the spontaneity of the reaction because it represents the result of two factors: the change in heat capacity ΔH and the factor of change in the entropy of the reaction ΔS [27], and the supply capacity of any soil can be found from the energy levels of the nutrients in the soil solution.

The level of this energy can represent the total differential of the supply forces of the element, which includes the nature of the ion, its salts, and its quality in relation to certain sites, as well as the ions accompanying it. It is possible to calculate the free energy from the following equation (ΔG = -RT InAR⁰) proposed by Woodruff [28] based on evaluate the ionic activity ratios (AR⁰) at equilibrium for potassium when no gain or loss occurs. A negative value for the amount of change in energy means that the reaction is spontaneous and does not require heat, while a positive value means that the reaction is nonspontaneous and requires heat. Through the results we obtained, it was shown that the values of the change in free energy ΔG for potassium in

the study soil were all negative, which this means that the reaction is spontaneous.

Table 6 displays the results for free energy, from which it is noted that the values of change in the free energy of potassium ΔG_{K} in the soil ranged from (-14.855×10⁻³ kJ) in the subsurface depth of the first site to (-11.724×10⁻³ kJ) in the surface depth of the second site at a rate of (-13.52×10⁻³ kJ) at a temperature of 278° K, While the values ranged from (-13.707×10⁻³ kJ) in the subsurface depth of the third site to $(-10.568 \times 10^{-3} \text{ kJ})$ in the subsurface depth of the first site, and the average reached (-12.133×10⁻³ kJ), at a temperature of 298° K. As the temperature increased to 318° K, the values ranged from (-14.821×10⁻³ kJ) at the subsurface depth of the second site to (-12.117×10⁻³ kJ) at the surface depth of the first site, with an average of (-13.19×10⁻³ kJ).

This difference in the free energy values of the study soils is directly related to the equilibrium conditions between the soil solution and its solid phase, as well as to the difference in exchange sites and its mineral composition [29]. Negative values of free energy indicate a high available of the potassium ion, and the higher the value with

Table 6. Value of free energy at equilibrium (ΔG) for three temperatures

Depth (cm)	(Cal	/ mole ⁻¹) at temperature	e Free energy (10 ⁻³)	
	278° Kelvin	298° Kelvin	318° Kelvin	
First Site				
Surface (0-25)	-14.342	-10.821	-12.117	
Sub Sur. (25-50)	-14.855	-10.568	-12.170	
Second Site				
Surface (0-25)	-11.724	-10.965	-12.663	
Sub Sur. (25-50)	-14.037	-13.681	-14.821	
Third Site				
Surface (0-25)	-13.424	-13.071	-13.480	
Sub Sur. (25-50)	-12.762	-13.707	-13.898	

a negative sign, the greater the availability of potassium quantitatively in the soil. Conversely, the greater the values of free energy with a positive sign mean a decrease in the available of potassium in those soils, which means the soil needs to add potassium [27].

3.5 Effect of Temperature on the Entropy of Potassium Adsorption Reactions (ΔS_{κ})

The entropy of the reaction (ΔS) indicates the regularity and arrangement of the ions and is one of the thermodynamic functions, and the change in its value is a sensitive measure of any change in the structure of the system [16]. It was observed in Table 7 that the increase in temperature led to making the interaction entropy values for all studied soils and for both depths less negative (higher reaction entropy values). which indicates the effect of temperature on the irregularity of potassium ions in the soil system and making them more entropy due to their higher internal energy due to the absorption of heat from the external environment affecting the reaction medium. The values of the entropy of the reaction at a temperature of 278 ranged from the highest value -53.43 expressed in joules.mol-¹ at the subsurface depth of the first site to the lowest value of -42.17 at the surface depth of the second site. While the increase in temperature to 298° led to an increase in the entropy values of the reaction system, the highest value of reaction entropy was recorded -45.99 in the subsurface depth of the third site and the lowest value of -35.46 in the subsurface depth of the first site. At 318' degrees, the highest value was -46.6 at the subsurface depth of the second site, and the lowest value was -38.10 at the surface depth of the first site. The change in the values of ΔS and its effect on the solubility of potassium-bearing minerals and its release from the solid soil phase

and its transfer to the soil solution will cause a change in the arrangement of the reaction system. For example, when the value of the change in ΔS is negative, it indicates the spontaneity of the reaction (that is, the reaction It occurs naturally and does not require a catalyst and is a heat emitter) and the decrease in negativity of ΔS values will be with increasing temperature [30].

3.6 Effect of Temperature on Enthalpy of Potassium Adsorption

Enthalpy refers to the heat released or absorbed during the process of dissolution or release and other reactions that occur in soil systems, and the change in enthalpy of the adsorption process is a measure of the heat of reaction. The results shown in Table 8 show that all enthalpy values for the study samples were negative values, which indicates that the reaction is of the exothermic type, as when the temperature increases, the enthalpy values increase (the negativity of the values decreases). The highest negative value for enthalpy was observed at the temperature of 278° K was -29.70, expressed in kJ.mol⁻¹, at the subsurface depth of the first site, and the lowest negative value was -23.44, at the surface depth of the second site. When the temperature rose to 298° K, the highest value of enthalpy was -26.49 at the subsurface depth in the third location, and the lowest value was -20.42 in the subsurface depth in the first location, while the highest value of enthalpy at a temperature of 318° K reached -27.77 in the subsurface depth, for the second site, compared to the lowest value -22.70 at the surface depth for the first site. The high values of enthalpy of the potassium adsorption process from the liquid soil phase indicate that it may be due to the partial or complete removal of the aqueous shells of potassium, calcium, and magnesium ions [31].

Table 7. Entropy values for adsorption reactions (ΔS_{K}) for three temperation

Depth (cm)	Entropy of reaction (joule.mol ⁻¹) at temperature					
	278° Kelvin	298° Kelvin	318° Kelvin			
First Site						
Surface (0-25)	-51.59	-36.31	-38.10			
Sub Sur. (25-50)	-53.43	-35.46	-38.27			
Second Site						
Surface (0-25)	-42.17	-36.79	-39.82			
Sub Sur. (25-50)	-50.49	-45.90	-46.60			
Third Site						
Surface (0-25)	-48.28	-43.86	-42.39			
Sub Sur. (25-50)	-45.90	-45.99	-43.70			

Al-Jubory and Al-Khafagi; Asian J. Soil Sci. Plant Nutri., vol. 10, no. 4, pp. 604-614, 2024; Article no.AJSSPN.127728

Depth	Enthalpy of reaction (K joule.mol ⁻¹) at temperature					
	278° Kelvin	298° Kelvin	318° Kelvin			
First Site						
Surface (0-25)	-28.68	-20.91	-22.70			
Sub Sur. (25-50)	-29.70	-20.42	-22.80			
Second Site						
Surface (0-25)	-23.44	-21.19	-23.73			
Sub Sur. (25-50)	-28.07	-26.44	-27.77			
Third Site						
Surface (0-25)	-26.84	-25.26	-25.26			
Sub Sur. (25-50)	-25.52	-26.49	-26.04			

Table 8. Enthalpy values for adsorption reactions (ΔH_K) for three temperatures

4. CONCLUSION

The ability to release potassium from the soil may be limited due to the change in the structure of potassium-bearing soil minerals and the decrease in the diffusion of potassium ions in the interlayer spaces of the minerals. The use of thermodynamic criteria in evaluating soil fertility to express potassium status and behavior in the soil helps in revealing an appropriate management method for each soil.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Authors hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- 1. Bayraktar SN. The role of potassium fertilizer and gypsum content in the soil in alleviating salt stress resulting from salinity of irrigation water and growth and yield of wheat. Master's thesis, College of Agriculture, Tikrit University; 2019. Iraq.
- 2. Mouhamad R, Alsaede A, Iqbal M. Behavior of potassium in soil: A mini review. Chem Int. 2016;2(1):58–69.
- Al-Obaidi MAJ, Al-Zubaidi AH. The effect of temperature on the adsorption and release of potassium in soil. Mesopotamian Agric J. 2005;33(4):47–56.
- 4. Al-Obaidi MAJ, Jamal RI, Abdullah FAH. The effect of wetting and drying cycles on the release of potassium added to some

calcareous soils from northern Iraq. Al-Rafidain Agric J. 2010;38(1):1–13.

- 5. Al-Samarrai OA. Status and behavior of potassium in protected agricultural soils. Doctoral thesis, College of Agriculture, University of Baghdad; 2005. Iraq.
- 6. Al-Shibini JM. Potassium in the ground and plants. 1st ed. Egyptian Library for Printing, Publishing and Distribution. 2007;207.
- Najafi-Ghiri M, Boostani HR, Mirsoleimani A, Mohaviye-Asadi N, Beizavi M, Shafiei M, et al. Potassium fixation and release in some calcareous soils under orange cultivation. Eurasian Soil Sci. 2020;53: 978–985.
- Isnaini S, Rivaie AA. Comparison of potassium quantity-intensity relationships in tropical paddy soil under tillage and notillage systems after fifteen growing seasons. Plant Soil Environ. 2023;69(1):1– 9.
- 9. Rani K, Datta A, Jat HS, Choudhary M, Sharma PC, Jat ML. Assessing the availability of potassium and its quantityintensity relations under long term conservation agriculture-based cereal systems in North-West India. Soil Tillage Res. 2023;228:105644.
- Carter MR, Gregorich EG. Soil sampling and methods of analysis. 2nd ed. Lewis Publishers, Canadian Society of Soil Science; 2008. CRC Press.
- Beckett PHT. Studies on soil potassium: confirmation of the ratio law: I. measurement of potassium potential. II. The immediate Q/I relation of labile potassium in soil. J Soil Sci. 1964;15:1– 23.
- 12. Moharami S, Jalali M. Effects of cations and anions on iron and manganese sorption and desorption capacity in calcareous soils from Iran. Environ Earth Sci. 2013;68(3):847–858.

Al-Jubory and Al-Khafagi; Asian J. Soil Sci. Plant Nutri., vol. 10, no. 4, pp. 604-614, 2024; Article no.AJSSPN.127728

- Ogawada RA, Sparks DL. A critical evaluation on the use of kinetics for determining thermodynamics of ion exchange in soils. Soil Sci Soc Am J. 1986;50:300–305.
- 14. Thajeel AS. Isotherm, Kinetic and Thermodynamic of adsorption of heavy metal ions onto local activated carbon. Aquat Sci Technol. 2013;1(2):1–9.
- 15. Al-Qattan RA. Impact of Wetting, Drying and Soil Texture on Dynamic and Kinetic Equilibrium for Cations in Some Soils of Nineveh Governorate. Master's thesis, College of Agriculture and Forestry, University of Mosul; 2023.
- 16. Sparks DL, Singh B, Siebecker MG. Environmental soil chemistry. 3rd ed. Elsevier; 2022.
- 17. Schindler FV, Howard J, Doolittle J. Assessment of soil potassium sufficiency as related to quantity-intensity in montmorillonitic soils. Commun Soil Sci Plant Anal. 2005;36:2255–2270.
- Bhonsle NS, Kumar S, Sekhon GS. Relationship of K forms and release characteristics with day mintilogy. Geoderma. 1992;54:285–293.
- 19. Sparks DL. Methods of soil analysis. Soil Sci Soc Am J. 2017;5585:1–13.
- 20. Al-Jubouri LM. Using some chemical standards and geographic information systems to evaluate the readiness of potassium and its relationship to the mineral composition in the soil of Babylon Governorate. Master's thesis, College of Agriculture, Al-Qasim Green University; 2017. Iraq.
- Li X, Xing Y, Tang L, Liu N, Chang Q, Zhang J. Adsorption of cations at the illite– water interface and its effect on intrinsic potassium ions. Eur J Soil Sci. 2022;73(1): e13155.
- 22. Das D, Dwivedi BS, Datta SP, Datta SC, Meena MC, Agarwal BK, et al. Potassium

supplying capacity of red soil from eastern India after forty-two years of continuous cropping and fertilization. Geoderma. 2019;341:76–92.

- 23. Wakeel A, Ishfaq M. Potash use and dynamics in agriculture. In: Wakeel A, editor. Potash use and dynamics in agriculture. Singapore: Springer. 2022;7–17.
- 24. Hamed MH, Amin AZ. Evaluation of potassium quantity-intensity in some soils of El-Dakhla Oasis, New Valley, Egypt. Alexandria Sci Exch J. 2017;38:112–119.
- 25. Bilias F, Barbayiannis N. Potassium availability: An approach using thermodynamic parameters derived from quantity-intensity relationships. Geoderma. 2019;338:353–364.
- 26. Strauss CR, Brummener GW. Influss der Kristallinität Van Phosphatadsorption. M: Heitungen der Deutschen Bodenkundlichen Gesellschaft; 1989.
- 27. Al-Obaidi MAJ, Fayyadh MA, Semo A. Kinetics of potassium adsorption and desorption in some Vertisols of Duhok Governorate, Kurdistan Region-Iraq; 2015.
- 28. Woodruff CM. Ionic equilibrium between clay and dilute salt solution; 1955.
- 29. Mahmmadani LA. Study of the effect of carbonate minerals and iron oxides removal on potassium adsorptiondesorption from clay fraction in some soils of northern Iraq. Doctoral thesis, College of Agriculture, University of Mosul; 2008.
- Prasad B, Sarangthem I. Thermodynamics of zinc-chelate adsorption in calcareous soils. J Indian Soc Soil Sci. 1992;40:458– 463.
- Apple C, Rhue RD, Ma LQ, Reve W. Enthalpies of K/Ca and K/Pb exchange in two tropical soils as measured by flow calorimetry. Soil Sci. 2002;167:773– 781.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history: The peer review history for this paper can be accessed here: https://www.sdiarticle5.com/review-history/127728