



## Mathematical Model of the Transmission Dynamics of Lassa Fever with Separation of Infected Individual and Treatment as Control Measures

S. C. Nwasuka<sup>1\*</sup>, I. E. Nwachukwu<sup>1</sup> and P. C. Nwachukwu<sup>2</sup>

<sup>1</sup>Department of Mathematics and Statistics, Clifford University, Owerrinta, Abia State, Nigeria.

<sup>2</sup>Department of Biological Science, Clifford University, Owerrinta, Abia State, Nigeria.

### Authors' contributions

This work was carried out in collaboration among all authors. Author NSC formulated and analyzed the model. Obtained the stabilities and carried out sensitivity analysis and simulation analysis, author NIJ helped in the literature search and was involved in the simulation analysis, author NPC helped with all the Biological/Medical terms and their explanation as needed and also help in the literature search. All the authors took part in the formatting.

### Article Information

DOI: 10.9734/JAMCS/2019/v32i630164

#### Editor(s):

- (1) Dr. Morteza Seddighin, Professor, Indiana University East Richmond, USA.  
(2) Dr. José González Enriquez, Professor, Department of Computer Languages and Systems, University of Sevilla, Spain.

#### Reviewers:

- (1) Devipriya Ganesan, Bharathiar University, India.  
(2) Abdullah Sonmezoglu, Yozgat Bozok, Turkey.  
(3) Anthony Spiteri Staines, University of Malta, Malta.  
(4) Utku Kose, Suleyman Demirel University, Turkey.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/48982>

Received: 01 March 2019

Accepted: 15 May 2019

Published: 04 July 2019

Short Research Article

### Abstract

A non-linear deterministic mathematical model is formulated and analysed to study the controllability of lassa fever incorporating separation of infected individuals and treatment measures. The model assumes that humans susceptible acquired the Infection via interaction with the infected rodent populations at a constant rate and also the model assumes that treatment is only given to separated human population. The existence, uniqueness and positivity of the model's solution have been carried out and the results shows that the solution exist and is unique. Again, the disease – free equilibrium state was obtained and analysed. We obtained an important threshold parameter called the effective reproduction number  $\mathcal{R}_{eff}$  using the next generation method. If  $\mathcal{R}_{eff} < 1$  the disease-free equilibrium exists and is locally and globally asymptotically stable, implying that Lassa fever can be controlled and eradicated within the population in a finite time and if the  $\mathcal{R}_{eff} > 1$ , the disease invade and become endemic in the population.

\*Corresponding author: E-mail: [nwasukac@clifforduni.edu.ng](mailto:nwasukac@clifforduni.edu.ng);

*Keywords: Lassa fever; mathematical model; separation of infected human; treatment; existence.*

## 1 Introduction

Lassa Fever (LF), technically known as Lassa Hemorrhagic Fever (LHF) is a deadly infectious illness to man caused by a Lassa Virus (LASV) or Lassa Hemorrhagic Fever Virus (LASHFV) from a carrier "multimamm-mate rat" (Genus name *Mastomys natalensis*) [1]. These kind of rat (multimamm-mate rat) are found in abundant in the sub-saharan part of Africa and infected rodent with the virus serves as a reservoir or host to the lassa virus (LASV) and transport it within the region in West Africa and some areas beyond. According to many researchers [1,2,3,4,5], Lassa fever is an acute viral infection associated with a wide spectrum of disease manifestations, which range from mild to hemorrhagic fever characterized by multiorgan failure.

Lassa fever virus is mainly a zoonosis (a disease that is animal-borne or transmitted to humans from animals), specifically an African rat, also called the natal multimammate rodent (*Mastomys natalensis*) serves as a host or reservoir of the virus. Once the rat has become a carrier, it will excrete the virus throughout the rest of its lifetime through feces and urine creating ample opportunity for exposure [2,6,7,8]. The multimammate rat can quickly produce a large number of offspring, tends to colonize human settlements increasing the risk of rodent-human contact, and is found throughout the west, central and eastern parts of the African continent [2,9,10,11]. The virus is probably transmitted by contact with the faeces or urine of animals accessing grain stores in residences Werner et al. [3]. Transmission or Infection of Lassa virus to humans typically occurs by direct or indirect exposure to animal excrement through the respiratory or gastrointestinal tracts or eating contaminated food, touching soiled objects, or exposure to open cuts or sores [12,13,14,15,16]. Inhalation of tiny particles of infectious material (aerosol) is believed to be the most significant means of exposure. It is possible to acquire the infection through broken skin or mucous membranes that are directly exposed to infectious material Because *Mastomys* rodents often live in and around homes and scavenge on leftover human food items or poorly stored food. *Mastomys* rodents are sometimes consumed as a food source and infection may occur when rodents are caught and prepared. Direct contact with infected rodents is not the only way in which people are infected; person-to-person transmission may occur after exposure to virus in the blood, tissue, secretions, or excretions of a Lassa virus-infected individual, presenting a disease risk for healthcare workers (called nosocomial transmission) where proper personal protective equipment (PPE) is not available or not used [17,18,19]. Lassa virus may be spread in contaminated medical equipment, such as reused needles. Also, during sexual intercourse, the virus can be transmitted because the virus is present in urine for between three and nine weeks after infection, and it can be transmitted in semen for up to three months after becoming infected [4] and "Lassa fever" [5,19]. Casual contact (including skin- to-skin contact without exchange of body fluids) does not spread Lassa virus. Finally, No study has proven presence of lassa virus in the breast milk, but the high level of viremia suggests it may be possible, [6]. Above all, individuals who are at a higher risk of contracting the infection are those who live in rural areas where *Mastomys* are discovered, and where sanitation is not prevalent [20,21].

Lassa fever is endemic in parts of west Africa including Sierra Leone, Liberia, Guinea and Nigeria; however, other neighboring countries are also at risk, as the animal vector for Lassa virus, the "multimammate rat" (*Mastomys natalensis*) is distributed throughout the region. In 2009, the first case from Mali was reported in a traveler living in southern Mali; Ghana reported its first cases in late 2011. Separation cases have also been reported in Côte d'Ivoire and Burkina Faso and there is serologic evidence of Lassa virus infection in Togo and Benin. The number of Lassa virus infections per year in west Africa is estimated at 100,000 to 300,000, with approximately 5,000 deaths. Unfortunately, such estimates are crude, because surveillance for cases of the disease is not uniformly performed. In some areas of Sierra Leone and Liberia, it is known that 10%-16% of people admitted to hospitals every year have Lassa fever, which indicates the serious impact of the disease on the population of this region. In this paper, a non – linear deterministic mathematical model of Lassa fever shall be formulated to study the impact of Transmission Dynamics on Lassa Fever Incorporating separation and Treatment as a Control Measures. We wish to show that our results both analytical and numerical with the control measures can reduce the spread of Lassa fever to an optimal level in infinite time.

## 1.1 The specific objective of the study

The specific objective of these study are to:

1. To formulate and analyse a mathematical model on the controllability of lassa fever incorporating isolation and treatment measures in terms of the reproduction number.
2. To determine the stability of the equilibrium points.
3. To understand how separation and treatment can reduce mortality rate among the infected individuals.
4. To contribute on how separation can reduce the force of infection rate among the unaffected individuals.

## 1.2 Model formulation and analysis

### 1.2.1 Assumptions of the model

- i. The total human and rodent populations are given by  $N_H = S_H + E_H + I_H + J_H + T_H$  and  $N_R = S_R + I_R$  respectively (Table 1).
- ii. We assumed that treatment is only given to separate human population.
- iii. Recruitment into the susceptible population is either by birth or immigration.
- iv. Members of the infected human population can as well move to the susceptible human population via treatment.
- v. Infection is acquired via interaction unlike the direct contact in the existing model.

**Table 1. State variables and parameters of the model**

Variables/parameters	Description
$S_H$	Susceptible human at time $t$
$E_H$	Exposed human at time $t$
$I_H$	Infected human at time $t$
$J_H$	separated human at time $t$
$T_H$	Treated human at time $t$
$S_R$	Susceptible rodent at time $t$
$I_R$	Infected rodent at time $t$
$\pi_H$	Recruitment rate into the susceptible human
$\pi_R$	Recruitment rate into the susceptible rodent
$\mu_H$	Natural death rate in human
$\mu_R$	Natural death rate in rodent
$\beta_2$	Effective contact between infected human and susceptible human
$\beta_1$	Effective contact between infected rats and either susceptible human or susceptible rodent
$\alpha$	Progression rate to active Lassa fever
$\phi$	Separation rate
$\tau$	Treatment rate
$\omega$	Recovery rate
$\delta$	Human disease induced death
$\beta$	Probability of getting Lassa fever
$N_H$	Total population of human
$N_R$	Total population of rodent
$\lambda$	Force of infection




Fig. 1. Schematic diagram of our model

### 1.3 The model equations

From the above assumptions and the schematic diagram, the model will be governed by the following non – linear differential equation;

$$\frac{dS_H}{dt} = \pi_H + \omega T_H - \lambda S_H - \mu_H S_H \quad (1)$$

$$\frac{dE_H}{dt} = \lambda S_H - (\mu_H + \alpha) E_H \quad (2)$$

$$\frac{dI_H}{dt} = \alpha E_H - (\mu_H + \delta + \phi) I_H \quad (3)$$

$$\frac{dJ_H}{dt} = \phi I_H - (\mu_H + \delta + \tau) J_H \quad (4)$$

$$\frac{dT_H}{dt} = \tau J_H - (\mu_H + \omega) T_H \quad (5)$$

$$\frac{dS_R}{dt} = \pi_R - \beta_1 S_R I_R - \mu_R S_R \quad (6)$$

$$\frac{dS_R}{dt} = \beta_1 S_R I_R - \mu_R I_R \quad (7)$$

where

$$\lambda = \beta_1 I_R - \beta_2 I_H \quad (8)$$

With the initial conditions  $S_H(0) = S_H^0, E_H(0) = E_H^0, I_H(0) = I_H^0, J_H(0) = J_H^0, T_H(0) = T_H^0, S_R(0) = S_R^0, I_R(0) = I_R^0$  and  $N(0) = N_0$ . The force of infection  $\lambda = \beta_1 I_R - \beta_2 I_H$ , where  $\beta_1$  and  $\beta_2$  are the effective contact between infected rats and either susceptible humans or susceptible rodents and effective contact between infected human and susceptible human respectively with,

$$N_H(t) = S_H(t) + E_H(t) + I_H(t) + J_H(t) + T_H(t) \quad (9)$$

Where  $N_H(t)$  denotes the total human population at a given time with its time derivative given by;

$$\frac{dN_H}{dt} = \frac{dS_H}{dt} + \frac{dE_H}{dt} + \frac{dI_H}{dt} + \frac{dJ_H}{dt} + \frac{dT_H}{dt} \quad (10)$$

Plugging (1) – (5) into (10) gives

$$\frac{dN_H}{dt} = \pi_H - (I_H + J_H)\delta - \mu_H N_H \quad (11)$$

Also,

$$N_R(t) = S_R(t) + I_R(t) \quad (12)$$

Where  $N_R(t)$  denotes the total rodents population at a given time with its time derivative given by;

$$\frac{dN_R}{dt} = \frac{dS_R}{dt} + \frac{dI_R}{dt} \quad (13)$$

Substituting (6) and (7) into (13) gives

$$\frac{dN_R}{dt} = \pi_R - \mu_R N_R \quad (14)$$

#### 1.4 The invariant region (region of biological interest)

As the system (1) – (5) monitors human population, all related state variables and parameters are assumed to be non – negative for all  $t \geq 0$ . Therefore, the above system is dissipative in the proper subset  $\Gamma \subset \mathbb{R}_+^5$ . Thus, we state and prove the following results:

**Lemma 1:** The solutions of the system (1) – (7) are feasible for all  $t > 0$  if they enter the invariant region  $\Gamma = (S_H, E_H, I_H, J_H, T_H, S_R, I_R)$ .

**Proof 1:**

Let  $\Gamma = (S_H, E_H, I_H, J_H, T_H, S_R, I_R)$  be any solution of the system (1) – (7), with non – negative initial conditions. From equation (11), we see that in the absence of Lassa fever ( $I_H = J_H = 0$ ), we obtained;

$$\frac{dN_H}{dt} \leq \pi_H - \mu_H N_H \quad (15)$$

Rearranging (15) gives

$$\frac{dN_H}{dt} + \mu_H N_H \leq \pi_H \tag{16}$$

Solving (16) using the method of integrating factor (IF) we compute the IF as follows:

$$IF = e^{\int \mu_H dt} = e^{\mu_H t} \tag{17}$$

Multiplying both sides of (16) by (17) yields

$$e^{\mu_H t} \frac{dN_H}{dt} + \mu_H N_H e^{\mu_H t} \leq \pi_H e^{\mu_H t}$$

That is,

$$\frac{d}{dt} (\mu_H N_H e^{\mu_H t}) \leq \pi_H e^{\mu_H t} \tag{18}$$

Integrating both sides of (18) gives

$$N_H(t) e^{\mu_H t} \leq \frac{\pi_H}{\mu_H} e^{\mu_H t} + \psi \tag{19}$$

Where  $\psi$  is a constant of integration.

This means

$$N_H(t) \leq \frac{\pi_H}{\mu_H} + \psi e^{-\mu_H t} \tag{20}$$

Applying the initial condition:  $N_H(0) = N_H^0$ , we obtain;

$$N_H^0 - \frac{\pi_H}{\mu_H} \leq \psi \tag{21}$$

Substituting (21) into (20) we have

$$N_H(t) \leq \frac{\pi_H}{\mu_H} + \left( N_H^0 - \frac{\pi_H}{\mu_H} \right) e^{-\mu_H t}$$

Applying Birkhoff and Rota's theorem on differential inequality (Birkhoff and Rota, 1982), we have  $0 \leq N_H(t) \leq \frac{\pi_H}{\mu_H}$ , as  $t \rightarrow \infty$ .

The total population approaches  $K = \frac{\pi_H}{\mu_H}$ , as  $t \rightarrow \infty$ . which is commonly known as the carrying capacity. Therefore, the feasible solutions set of the extended model (1) – (5) enters the region below

$$\left\{ \Gamma = \Gamma = (S_H, E_H, I_H, J_H, T_H) \in \mathbb{R}_+^5 : S_H > 0, E_H \geq 0, I_H \geq 0, J_H \geq 0, T_H \geq 0, N_H \leq \frac{\pi_H}{\mu_H} \right\}.$$

Thus in this region our model is biologically feasible. Here whenever  $N > \frac{\pi_H}{\mu_H}$  then  $\frac{dN_H}{dt} < 0$  which means the population reduces asymptotically to the carrying capacity and whenever  $N \leq \frac{\pi_H}{\mu_H}$  every solution with initial condition in  $\Gamma$  remains in that region for  $t > 0$ , so the model is well posed in  $\Gamma$ . Therefore, the region  $\Gamma$  is positively – invariant (i.e. solutions remain positive for all time.) and the model is well posed and biologically meaningful and this ends the proof of the Lemma 1.

### 1.5 Existence of disease free equilibrium state ( $\mathcal{E}_0$ ) of the model

Here, we compute the model disease free equilibrium state by setting the time – derivatives on the right hand sides of the model system (1) – (9) to zero such that

$$\frac{dS_H}{dt} = \frac{dE_H}{dt} = \frac{dI_H}{dt} = \frac{dJ_H}{dt} = \frac{dT_H}{dt} = \frac{dS_R}{dt} = \frac{dI_R}{dt} = 0 \quad (22)$$

So that we now have

$$0 = \pi_H + \omega T_H - (\beta_1 I_R + \beta_2 I_H) S_H - \mu_H S_H \quad (23)$$

$$0 = (\beta_1 I_R + \beta_2 I_H) S_H - (\mu_H + \alpha) E_H \quad (24)$$

$$0 = \alpha E_H - (\mu_H + \delta + \phi) I_H \quad (25)$$

$$0 = \phi I_H - (\mu_H + \delta + \tau) J_H \quad (26)$$

$$0 = \tau J_H - (\mu_H + \omega) T_H \quad (27)$$

$$0 = \pi_R - \beta_1 S_R I_R - \mu_R S_R \quad (28)$$

$$0 = \beta_1 S_R I_R - \mu_R I_R \quad (29)$$

Recall that, the disease free equilibrium state of the model (1) – (9) is scenario where there is no disease in the system which implies that

$$E_H = I_H = J_H = T_H = I_R = 0$$

Plugging (31) into (1) – (9) and solving accordingly we obtain

$$\mathcal{E}_0 = (S_H^0, E_H^0, I_H^0, J_H^0, T_H^0, S_R^0, I_R^0) = \left( \frac{\pi_H}{\mu_H}, 0, 0, 0, 0, \frac{\pi_R}{\mu_R}, 0 \right) \quad (30)$$

## 1.6 Effective Reproduction Number ( $\mathcal{R}_{eff}$ )

$$\mathcal{R}_0 = \rho(FV^{-1})$$

where  $\rho(FV^{-1})$  is the spectral radius of next generation matrix.

We calculate the basic reproduction number using the next generation operator method on the system (1) – (9) as follows;

The vector  $F_i$  of the rates of the new infection in compartment  $E_H, I_H, J_H, T_H$  and  $I_R$  is given by

$$F_i = \begin{pmatrix} (\beta_1 I_R + \beta_2 I_H) S_H \\ 0 \\ 0 \\ 0 \\ \beta_1 I_R S_R \end{pmatrix} \quad (31)$$

Also, the remaining transfer terms in compartment  $E_H, I_H, J_H, T_H$  and  $I_R$  is given by

$$V_i = \begin{pmatrix} (\mu_H + \alpha) E_H \\ (\mu_H + \delta + \phi) I_H - \alpha E_H \\ (\mu_H + \delta + \tau) J_H - \phi I_H \\ (\mu_H + \omega) E_H - \tau J_H \\ \mu_H I_R \end{pmatrix} \quad (32)$$

The matrix of partial derivative of  $F_i$  at the disease free equilibrium state at  $\mathcal{E}_0 = (S_H^0, 0, 0, 0, S_R^0, 0)$  is given by

$$F(\mathcal{E}_0) = \begin{pmatrix} 0 & \frac{\beta_2 \pi_H}{\mu_H} & 0 & 0 & \frac{\beta_1 \pi_H}{\mu_H} \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \frac{\beta_1 \pi_R}{\mu_R} \end{pmatrix} \quad (33)$$

Rewriting (33) yields

$$F(\mathcal{E}_0) = \begin{pmatrix} 0 & F_{12} & 0 & 0 & F_{15} \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & F_{55} \end{pmatrix} \quad (34)$$

where;

$$\begin{aligned} F_{12} &= \frac{\beta_2 \pi_H}{\mu_H} \\ F_{15} &= \frac{\beta_1 \pi_H}{\mu_H} \\ F_{55} &= \frac{\beta_1 \pi_R}{\mu_R} \end{aligned} \quad (35)$$

Also, the matrix of the partial derivatives of  $V_i$  at the disease free equilibrium state  $\mathcal{E}_0 = (S_H^0, 0, 0, 0, S_R^0, 0)$  is given by

$$V(\mathcal{E}_0) = \begin{pmatrix} (\mu_H + \alpha) & 0 & 0 & 0 & 0 \\ -\alpha & (\mu_H + \delta + \phi) & 0 & 0 & 0 \\ 0 & -\phi & (\mu_H + \delta + \tau) & 0 & 0 \\ 0 & 0 & -\tau & (\mu_H + \omega) & 0 \\ 0 & 0 & 0 & 0 & \mu_R \end{pmatrix} \quad (36)$$

Equation (36) can also be written as;

$$V(\mathcal{E}_0) = \begin{pmatrix} V_{11} & 0 & 0 & 0 & 0 \\ -V_{21} & V_{22} & 0 & 0 & 0 \\ 0 & -V_{32} & V_{33} & 0 & 0 \\ 0 & 0 & -V_{43} & V_{44} & 0 \\ 0 & 0 & 0 & 0 & V_{55} \end{pmatrix} \quad (37)$$

where;

$$\left. \begin{aligned} V_{11} &= \mu_H + \alpha & V_{55} &= \mu_R \\ V_{22} &= \mu_H + \delta + \phi & V_{21} &= \alpha \\ V_{33} &= \mu_H + \delta + \tau & V_{32} &= \phi \\ V_{44} &= \mu_H + \omega & V_{43} &= \tau \end{aligned} \right\} \quad (38)$$

Computing the inverse of (37) gives

$$V^{-1} = \begin{pmatrix} \frac{1}{V_{11}} & 0 & 0 & 0 & 0 \\ \frac{V_{21}}{V_{11}V_{22}} & \frac{1}{V_{22}} & 0 & 0 & 0 \\ 0 & \frac{V_{32}}{V_{22}V_{33}} & \frac{1}{V_{11}} & 0 & 0 \\ \frac{V_{21}V_{32}V_{43}}{V_{11}V_{22}V_{33}V_{44}} & \frac{V_{43}}{V_{22}V_{44}} & \frac{V_{43}}{V_{33}V_{44}} & \frac{1}{V_{44}} & 0 \\ 0 & 0 & 0 & 0 & \frac{1}{V_{55}} \end{pmatrix} \quad (39)$$

Rewriting (39) we have;

$$V^{-1} = \begin{pmatrix} A_{11} & 0 & 0 & 0 & 0 \\ A_{21} & A_{22} & 0 & 0 & 0 \\ 0 & A_{32} & A_{33} & 0 & 0 \\ A_{41} & A_{42} & A_{43} & A_{44} & 0 \\ 0 & 0 & 0 & 0 & A_{55} \end{pmatrix} \quad (40)$$

where;

$$A_{11} = \frac{1}{(\mu_H + \alpha)} \quad (41)$$

$$A_{21} = \frac{\alpha}{(\mu_H + \alpha)(\mu_H + \delta + \phi)} \quad (42)$$

$$A_{22} = \frac{\phi}{(\mu_H + \delta + \phi)} \quad (43)$$

$$A_{32} = \frac{\phi}{(\mu_H + \delta + \phi)(\mu_H + \delta + \tau)} \quad (44)$$

$$A_{33} = \frac{1}{(\mu_H + \delta + \tau)} \quad (45)$$

$$A_{41} = \frac{\alpha\phi\tau}{(\mu_H + \alpha)(\mu_H + \delta + \phi)(\mu_H + \delta + \tau)(\mu_H + \omega)} \quad (46)$$

$$A_{42} = \frac{\tau}{(\mu_H + \delta + \phi)(\mu_H + \omega)} \quad (47)$$

$$A_{43} = \frac{\tau}{(\mu_H + \delta + \phi)(\mu_H + \omega)} \quad (48)$$

$$A_{44} = \frac{1}{(\mu_H + \omega)} \quad (49)$$

$$A_{42} = \frac{1}{\mu_R} \quad (50)$$

To compute  $FV^{-1}$  we use (41) and (43) so that;

$$FV^{-1} = \begin{pmatrix} 0 & F_{12} & 0 & 0 & F_{15} \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & F_{55} \end{pmatrix} \begin{pmatrix} A_{11} & 0 & 0 & 0 & 0 \\ A_{21} & A_{22} & 0 & 0 & 0 \\ 0 & A_{32} & A_{33} & 0 & 0 \\ A_{41} & A_{42} & A_{43} & A_{44} & 0 \\ 0 & 0 & 0 & 0 & A_{55} \end{pmatrix}$$

$$FV^{-1} = \begin{pmatrix} 0 & F_{11}A_{12} & 0 & 0 & F_{11}F_{15} \\ 0 & F_{12}A_{12} & 0 & 0 & F_{12}A_{15} \\ 0 & 0 & 0 & 0 & 0 \\ 0 & F_{41}A_{12} & 0 & 0 & F_{41}A_{15} \\ 0 & 0 & 0 & 0 & F_{55}A_{55} \end{pmatrix} \quad (51)$$

It follows that the effective reproduction number  $\mathcal{R}_{eff}$  is computed by taking the spectral radius (dominant eigenvalue) of the matrix  $FV^{-1}$  using the characteristics equation 36 and 42

$$\det(FV^{-1} - \lambda_E I) = 0 \quad (52)$$

$$\begin{matrix} & & \text{or} & & \\ \left| \begin{matrix} -\lambda & F_{11}A_{12} & 0 & 0 & F_{11}F_{15} \\ 0 & F_{12}A_{12} - \lambda & 0 & 0 & F_{12}A_{15} \\ 0 & 0 & -\lambda & 0 & 0 \\ 0 & F_{41}A_{12} & 0 & -\lambda & F_{41}A_{15} \\ 0 & 0 & 0 & 0 & F_{55}A_{55} - \lambda \end{matrix} \right| & = & 0 \end{matrix} \quad (53)$$

Evaluating (56) accordingly gives;

$$\lambda_1 = \lambda_3 = \lambda_4 = 0 \quad (54)$$

and

$$\lambda_2, \lambda_5 = \max \left( \frac{\beta_2 \alpha \pi_H}{\mu_H(\mu_H + \alpha)(\mu_H + \delta + \phi)}, \frac{\beta_1 \pi_R}{\mu_R} \right) \quad (55)$$

Therefore, the largest (dominant) eigenvalue also known as the effective reproduction denoted by  $\mathcal{R}_{eff}$  is given by

$$\mathcal{R}_{eff} = \frac{\beta_2 \alpha \pi_H}{\mu_H(\mu_H + \alpha)(\mu_H + \delta + \phi)} \quad (56)$$

with

$\frac{1}{\mu_H}, \frac{1}{(\mu_H + \alpha)}$  and  $\frac{1}{(\mu_H + \delta + \phi)}$  which refers to per capital human mortality,

**Table 2. Parameters values for numerical and sensitivity analysis**

Parameters	Values	Sources
$\pi_H$	2000	[18]
$\pi_R$	500	[18]
$\beta_1$	0.2	[18]
$\beta_2$	0.2	Estimated
$\alpha$	0.003	[18]
$\phi$	0.2	Estimated
$\tau$	0.75	Assumed
$\mu_H$	0.02	[18]
$\mu_R$	0.02	[18]
$\delta$	0.1	Assumed
$\omega$	0.54	Assumed

**Biological Interpretation 1:** The biological meaning of the parameter components of the effective reproduction number are as follows:

$\left(\frac{\pi_H}{\mu_H}\right)$ : The carrying capacity for human population.

$\left(\frac{\alpha}{\mu_H+\alpha}\right)$ : The proportion of individuals from the exposed human that becomes infectious.

$\left(\frac{\beta_2}{\mu_H+\delta+\phi}\right)$ : The average number of susceptible human infected by a single human infectious.

### 1.7 Numerical simulations

Here, we carryout numerical simulation of the model (1) – (9) using the set of reasonable parameters and initial values given in Tables 2 and 3 and 4 whose sources are mainly from [24] as well as assumed values based on the literature of the disease in order to have more realistic simulation results.

**Table 3. Parameters values for numerical and sensitivity analysis**

Variables	Values	Sources
$S_H(t)$	10000	[19]
$E_H(t)$	3000	Assumed
$I_H(t)$	2000	[19]
$J_H(t)$	1500	Assumed
$T_H(t)$	600	[19]
$S_H(t)$	200	[19]
$I_R(t)$	125	[19]

**Table 4. Sensitivity indices of  $\mathcal{R}_{eff}$**

S/N	Parameter	Sensitivity index	Sign
1	$\beta_1$	1.000000000	+
2	$\pi_H$	1.000000000	+
3	$\alpha$	0.8695652173	+
4	$\delta$	0.312500000	-
5	$\phi$	0.625000000	-
6	$\mu_H$	1.9320652180	-

## 2 Numerical Results

In this sub – section, we presents the numerical results of the above experiment as follows:

### 2.1 Simulation results showing the trends of the state variables of the lassa fever model with separation rate for the infected class

The impact of separation rate  $\phi$  on Lassa fever is investigated. The result of this is given in Fig. 2. We observed that, an increase in the separation rate brings about a declined in the infected humans. This is true because as separation for treatment increases, infected human population on the order hand reduces. This also shows that, treatment of Lassa fever will go a long way in curtailing the disease through separation and treatment of the infected humans.




Fig. 2. Simulation results showing the effect of separation rate  $\phi$  on infected humans




Fig. 3. Simulation results showing the impact of treatment rate  $\tau$  on infected individuals

## 2.2 Simulation results showing the trends of the state variables of the lassa fever model with recovery rate due to treatment for; infected humans and treated humans

Numerical results depicted on Fig. 4 indicates that, increasing treatment rate reduces the infected humans population, this is in line with reality because treatment of the infected persons through separation of infected individuals brings about reduction in number of person infected with Lassa fever.




Fig. 4. Simulation results showing the effect of treatment rate  $\tau$  on treated humans

## 3 Conclusion

Nigeria is endemic to Lassa Fever and has being rated as one of the country in West Africa with the high transmission rate of Lassa fever, but we should not panic as the impact of this paper will highly contribute in curbing Lassa fever since our Reproduction number is less than one ( $\mathcal{R}_{eff} < 1$ ), which simply implies that Lassa Fever can be eradicated from the country. Therefore we conclude that since our  $\mathcal{R}_{eff} < 1$  is less than one the disease will surely die out in infinite time.

## Competing Interests

Authors have declared that no competing interests exist.

## References

- [1] David Safronetz, Job E. Lopez, Nafomon Sogoba, Sékou F. Traore, Sandra J. Raffel, Elizabeth R. Fischer, Hideki Ebihara, Luis Branco, Robert F. Garry, Tom G. Schwan, Heinz Feldmann. Detection of lassa virus, mali article in emerging infectious diseases; 2010.  
DOI: 10.3201/eid1607.100146

- [Source: PubMed 21 May 2014]
- [2] Go AS, Bauman M, King SM, Fonarow GC, Lawrence W, Williams KA, Sanchez E. An effective approach to high blood pressure control: A science advisory from the American Heart Association, the American College of Cardiology, and the Centers for Disease Control and Prevention. *Hypertension*. 2013;63(4):878–85.  
DOI: 10.1161/HYP.0000000000000003  
[PMID 24243703]  
(Archived from the original on 20 November 2013)
- [3] Werner Dietrich. editor *Biological Resources and Migration*. Springer. 2004;363.  
[ISBN 978-3-540-21470-0]
- [4] Public Health England: Lassa fever: Origins, reservoirs, transmission and guidelines Archived 2 February 2016 at the Wayback Machine. First published: 5 September 2014. Last updated: 1 April 2016.
- [5] "Lassa fever". Media Centre Fact Sheet No 179. World Health Organization. Archived from the original on 5 June 2015.  
[Retrieved 26 May 2015]
- [6] David Greenky, Barbara Knust, Eric J. dziubanwhat pediatricians should know about lassa virus. *JAMA Pediatr*. 2018;172(5):407-408.  
DOI: 10.1001/jamapediatrics.2017.5223
- [7] Ogbua O, Ajuluchukwub E, Unekec CJ. Lassa fever in West African sub-region: An overview. *J Vect Borne Dis*. 2007;44:1–11.
- [8] Nadezhda E. Yun, David H. Walker. Pathogenesis of lassa fever. *Viruses*. 2012;4(10):2031-48.  
DOI: 10.3390/v4102031  
[Review.PMID:23202452]  
Available:<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3497040/>
- [9] Dimie Ogoina. Lassa fever: A clinical and epidemiological review article 2013. *Niger Delta Journal of Medicine and Medical Research*. 2013;1(1):1-10.  
Available:<https://www.researchgate.net/publication/276270028>, page was uploaded on 14 May 2015.
- [10] Centers for Disease Control and Prevention, "Lassa Fever" Archived 23 September 2016 at the Wayback Machine.  
[Retrieved 15 August 2018]
- [11] World Health Organization (WHO) Lassa Fever Outbreak In Nigeria; 2018.  
Available:<http://www.who.int/medicalcentre> 20 April 2018
- [12] Amorosa V, MacNeil A, McConnell R, et al. Imported lassa fever, Pennsylvania, USA; 2010. *Emerging Infectious Diseases*. 2010;16(10):1598-600.
- [13] Nigerian LASSA Fever Resurgence: Prevention & Control; 2018.  
Available:<http://russellsmithgroup.com/health-and-safety/nigerian-lassa-fever-resurgence-prevention-control/>, January 31<sup>st</sup> 2018
- [14] World Health Organization (WHO). Lassa Fever – Nigeria Emergencies preparedness, response. Available:<http://www.who.int/medicalcentre>, July 5<sup>th</sup> 2018

- [15] Okuonghae D, Okuonghae R. A mathematical model for lassa fever. Journal of the Nigerian Association of Mathematical Physics. 2006;10:457-464.
- [16] Obabiyi OS, Akindele A. Onifade. Global stability analysis for lassafever transmission dynamics with optimal control application. Article in International Journal of Applied Mathematics. 2018;31(3):457-481.  
[ISSN: 1311-1728 (printed version)]  
[ISSN: 1314-8060 (on-line version)]  
Available:<http://dx.doi.org/10.12732/ijam.v31i3.11> January 2018
- [17] Akanni JO, Adediipo AD. Sensitivity analysis of the dynamical transmission and control of lassa fever virus. Asian Research Journal of Mathematics. 2018;9(3):1-11.  
[Article: no.ARJOM.37441]  
[ISSN: 2456-477X]  
[Published: 20th April 2018]
- [18] Adewale SO, Olopade IA, Ajao SO, Adeniran GA, Oyedemi OT. Mathematical analysis of lassa fever model with isolation. Asian Journal of Natural & Applied Sciences; 2016.
- [19] Akanni JO, Adediipo AD. Sensitivity analysis of the dynamics transmission and control of lassa fever virus. Asian Research Journal of Mathematics. 2018;9(3):1-11.
- [20] Nwasuka SC, Nwala KT. Global stability for cholera epidemic model in Nigeria. International Digital Organization for Scientific Research ISSN: 2550-7931 Idosr Journal of Applied Sciences. 2018;3(3):1-13.
- [21] Osiogogu UA, Nwasuka SC, Efor TE. Modeling the effect of treatment, vaccination and public health education campaign on the transmission tuberculosis infectious. Interdisciplinary Science Reviews. 2019;14(1):70-85.

---

© 2019 Nwasuka et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Peer-review history:**

The peer review history for this paper can be accessed here (Please copy paste the total link in your browser address bar)  
<http://www.sdiarticle3.com/review-history/48982>